

## **ISOVUE® (Iopamidol Injection)**

### **INDICATION:**

**ISOVUE®-200, -250, -300, -370 (Iopamidol Injection):** is indicated for:

- angiography in adults throughout the cardiovascular system including cerebral and peripheral arteriography, coronary arteriography and ventriculography, selective visceral arteriography and aortography
- in pediatric patients for angiocardiology
- in adult and pediatric intravenous excretory urography
- contrast enhancement of computed tomographic (CECT) head and body imaging
- peripheral venography in adults
- Oral procedures in pediatric patients and adults for:
  - CT of the abdomen and pelvis to delineate the gastrointestinal tract in adults and pediatric patients

### **WARNING: RISKS ASSOCIATED WITH INTRATHECAL ADMINISTRATION**

**Intrathecal administration, even if inadvertent, can cause death, convulsions, cerebral hemorrhage, coma, paralysis, arachnoiditis, acute renal failure, cardiac arrest, seizures, rhabdomyolysis, hyperthermia, and brain edema. ISOVUE is for intra-arterial or intravenous use only.**

### **IMPORTANT SAFETY INFORMATION:**

#### **WARNINGS AND PRECAUTIONS**

##### **Risks Associated with Intrathecal Administration**

Intrathecal administration, even if inadvertent, can cause death, convulsions, cerebral hemorrhage, coma, paralysis, arachnoiditis, acute renal failure, cardiac arrest, seizures, rhabdomyolysis, hyperthermia, and brain edema. ISOVUE is for intra-arterial or intravenous use only and must not be administered intrathecally.

##### **Hypersensitivity Reactions**

ISOVUE can cause life-threatening or fatal hypersensitivity reactions including anaphylaxis. Manifestations include respiratory arrest, laryngospasm, bronchospasm, angioedema, and shock. Most severe reactions develop shortly after the start of injection (e.g., within 1 to 3 minutes), but delayed reactions can also occur. There is increased risk of hypersensitivity reactions in patients with a history of previous reactions to contrast agents, and allergic disorders (i.e., bronchial asthma, allergic rhinitis, and food allergies) or other hypersensitivities. Obtain a history of allergy, hypersensitivity, or hypersensitivity reactions to iodinated contrast agents and always have emergency resuscitation equipment and trained personnel available prior to ISOVUE administration. Monitor all patients for hypersensitivity reactions.

##### **Acute Kidney Injury**

Acute kidney injury, including renal failure, may occur after ISOVUE administration. Risk factors include: pre-existing renal insufficiency, dehydration, diabetes mellitus, congestive heart failure, advanced vascular disease, elderly age, concomitant use of nephrotoxic or diuretic medications, multiple myeloma or other paraproteinemias, and repetitive or large doses of ISOVUE. Use the lowest necessary dose of ISOVUE in patients with renal impairment.

Adequately hydrate patients prior to and following ISOVUE administration. Do not use laxatives, diuretics, or preparatory dehydration prior to ISOVUE administration.

### **Cardiovascular Adverse Reactions**

ISOVUE increases the circulatory osmotic load and may induce acute or delayed hemodynamic disturbances in patients with congestive heart failure, severely impaired renal function, combined renal and hepatic disease, and combined renal and cardiac disease, particularly when repetitive or large doses are administered. Fatal cardiovascular reactions have occurred mostly within 10 minutes of ISOVUE injection; the main feature was cardiac arrest with cardiovascular disease as the main underlying factor. Hypotensive collapse and shock have occurred. Cardiac decompensation, serious arrhythmias, and myocardial ischemia or infarction can occur during coronary arteriography and ventriculography. Use the lowest necessary dose of ISOVUE in patients with congestive heart failure and always have emergency resuscitation equipment and trained personnel available. Monitor all patients for severe cardiovascular reactions.

### **Thromboembolic Events**

Serious, in some cases fatal, thromboembolic events, including myocardial infarction and stroke, can occur during angiographic procedures. During these procedures, increased thrombosis and activation of the complement system occurs. Risk factors for developing thromboembolic events include: length of procedure, catheter and syringe material, underlying disease state, and concomitant medications. To minimize thromboembolic events, use meticulous angiographic techniques and minimize the length of the procedure. Avoid blood remaining in contact with syringes containing iodinated contrast agents, which increases risk of clotting. Avoid angiocardiology in patients with homocystinuria because of the risk of inducing thrombosis and embolism.

### **Extravasation and Injection Site Reactions**

Extravasation can occur with ISOVUE administration, particularly in patients with severe arterial or venous disease. Inflammation, blistering, skin necrosis, and compartment syndrome have been reported following extravasation. In addition, injection site reactions such as pain and swelling at the injection site can also occur. Ensure intravascular placement of catheters prior to injection. Monitor patients for extravasation and advise patients to seek medical care for progression of symptoms.

### **Thyroid Storm in Patients with Hyperthyroidism**

Thyroid storm has occurred after the intravascular use of iodinated agents in patients with hyperthyroidism or with an autonomously functioning thyroid nodule. Evaluate the risk in such patients before use of ISOVUE.

### **Thyroid Dysfunction in Pediatric Patients 0 Years to 3 Years of Age**

Thyroid dysfunction characterized by hypothyroidism or transient thyroid suppression has been reported after both single exposure and multiple exposures to iodinated contrast agents in pediatric patients 0 years to 3 years of age. Younger age, very low birth weight, prematurity, underlying medical conditions affecting thyroid function, admission to neonatal or pediatric intensive care units, and congenital cardiac conditions are associated with an increased risk of hypothyroidism after iodinated contrast agent exposure. Pediatric patients with congenital cardiac conditions may be at greatest risk given that they often require high doses of contrast during invasive cardiac procedures. An underactive thyroid during early life may be harmful for cognitive and neurological development and may require thyroid hormone replacement therapy.

After exposure to iodinated contrast agents, individualize thyroid function monitoring based on underlying risk factors, especially in term and preterm neonates.

### **Hypertensive Crisis in Patients with Pheochromocytoma**

Hypertensive crisis in patients with pheochromocytoma has occurred with iodinated contrast agents. Closely monitor patients when administering ISOVUE if pheochromocytoma or catecholamine-secreting<sup>7</sup> of 13 paragangliomas are suspected. Inject the minimum amount of ISOVUE necessary and have measures for treatment of hypertensive crisis readily available.

### **Sickle Cell Crisis in Patients with Sickle Cell Disease**

Iodinated contrast agents can promote sickling in individuals who are homozygous for sickle cell disease. Hydrate patients prior to and following ISOVUE administration and use only if the necessary imaging information cannot be obtained with alternative imaging modalities.

### **Severe Cutaneous Adverse Reactions**

Severe cutaneous adverse reactions (SCAR) may develop from 1 hour to several weeks after intravascular contrast agent administration. These reactions include Stevens-Johnson syndrome and toxic epidermal necrolysis (SJS/TEN), acute generalized exanthematous pustulosis (AGEP), and drug reaction with eosinophilia and systemic symptoms (DRESS). Reaction severity may increase and time to onset may decrease with repeat administration of a contrast agent; prophylactic medications may not prevent or mitigate severe cutaneous adverse reactions. Avoid administering ISOVUE to patients with a history of a severe cutaneous adverse reaction to ISOVUE.

### **Interference with Laboratory Tests**

ISOVUE can interfere with protein-bound iodine test.

## **ADVERSE REACTIONS**

The most frequent adverse reactions are pain, hot flashes, burning sensation, nausea, and warmth.

## **DRUG INTERACTIONS**

**Metformin:** In patients with renal impairment, metformin can cause lactic acidosis. Iodinated contrast agents appear to increase the risk of metformin-induced lactic acidosis, possibly as a result of worsening renal function.

**Radioactive Iodine:** Administration of ISOVUE may interfere with thyroid uptake of radioactive iodine (I-131 and I-123) and decrease therapeutic and diagnostic efficacy. Avoid thyroid therapy or testing for up to 6 weeks post ISOVUE.

## **USE IN SPECIFIC POPULATIONS**

**Geriatric Use:** Iopamidol is excreted by the kidney, and the risk of adverse reactions to ISOVUE may be greater in patients with renal impairment. Because patients 65 years of age and older are more likely to have renal impairment, care should be taken in dose selection, and it may be useful to monitor renal function.

**Renal Impairment:** The clearance of iopamidol decreases with increasing degree of renal impairment and results in delayed opacification of the urinary system. In addition, preexisting renal impairment increases the risk for acute kidney injury. Iopamidol can be removed by dialysis.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

**Please click [here](#) for full Prescribing Information for ISOVUE® products.**

ISOVUE is currently manufactured for Bracco Diagnostics Inc. at three locations: BIPSO GmbH, Singen (Germany), Patheon Italia S.p.A., Ferentino (Italy), and S. M. Farmaceutici SRL, Tito (Italy).

ISOVUE is a registered trademark of Bracco Diagnostics Inc.

Bracco Diagnostics Inc.  
510 Carnegie Center  
Suite 300  
Princeton, NJ 08540 USA  
Phone: 609-514-2200  
Toll-Free: 1-800-631-5245 (U.S. only)  
Fax: 609-514-2424

© 2026 Bracco Diagnostics Inc. All Rights Reserved.