E-Z-GAS II- antacid/antiflatulent granule, effervescent E-Z-EM, INC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

E-Z-GAS® II EFFERVESCENT GRANULES ANTACID/ANTIFLATULENT

Drug Facts

Active Ingredients (in each packet)	Purpose
Citric Acid 1.53 g	Antacid
Simethicone 0.04 g	Antiflatulent
Sodium Bicarbonate 2.21 g	Antacid

Uses

- relieves
 - heartburn acid indigestion
 - sour stomach upset stomach
- relieves gas associated with these symptoms

Warnings

Do not use

- In cases of known or suspected esophageal obstruction.
- To dislodge foreign bodies.
- More than the maximum recommended dosage in 24-hour period.
- the maximum dosage of this product for more than two weeks, except under the advice and supervision of a physician.

Ask a doctor before use

- if you are taking a prescription drug. Antacids may interact with certain prescription drugs.
- If you have a sodium restricted diet.

Keep out of reach of children. In case of overdose seek medical help or contact a poison center immediately.

Once package is opened, discard unused portion.

Directions

	Recommended dose	Maximum recommended dose
Adults (under 60 years)	One packet every 6 hours or as directed by physician	Not to exceed 4 packets in 24 hours
Adults (60 years and older)	One packet every 8 hours or as directed by physician	Not to exceed 3 packets in 24 hours
Children (6 to 12 years)	One packet every 8 hours or as directed by physician	Not to exceed 3 packets in 24 hours
Children Under 6 years	Consult physician	Consult physician

Other information

- Store at controlled room temperature 15° 30° C (59° 86°F).
- Protect from moisture.
- Sodium content: approximately 26 mEq (600 mg) per dose.

Inactive ingredients

Maltodextrin, natural orange flavor, sodium saccharin. Questions? 1-516-333-8230 or 1-800-5444624

E-Z-Gas II

NDC: 10361-793-01

00105D Rev 10/2014